

2009

## The Fate of Potassium Ions Released from Contractile Muscle during Repeated Supramaximal Exercise

Anna Szczesna-Kaczmarek

*Department of Physiology, Jędrzej Sniadecki Academy of Physical Education and Sport in Gdansk, Poland, annaszka@awf.gda.pl*

Follow this and additional works at: <https://dcdgansk.bepress.com/journal>



Part of the [Health and Physical Education Commons](#), [Sports Medicine Commons](#), [Sports Sciences Commons](#), and the [Sports Studies Commons](#)

---

### Recommended Citation

Szczesna-Kaczmarek A. The Fate of Potassium Ions Released from Contractile Muscle during Repeated Supramaximal Exercise. *Balt J Health Phys Act.* 2009;1:20-26. doi: 10.2478/v10131-009-0002-0

This Article is brought to you for free and open access by Baltic Journal of Health and Physical Activity. It has been accepted for inclusion in Baltic Journal of Health and Physical Activity by an authorized editor of Baltic Journal of Health and Physical Activity.

## The Fate of Potassium Ions Released from Contractile Muscle during Repeated Supramaximal Exercise

DOI: 10.2478/v10131-009-0002-0

### Authors' Contribution:

A – Study Design  
B – Data Collection  
C – Statistical Analysis  
D – Data Interpretation  
E – Manuscript Preparation  
F – Literature Search  
G – Funds Collection

**Anna Szczęśna-Kaczmarek**

Jedrzej Sniadecki Academy of Physical Education and Sport in Gdansk,  
Poland

**Key words:** *supramaximal exercise, potassium ion, acidification, blood compartments, renal secretion*

### Abstract

**Background:** *The purpose of this paper was to examine the fate of K<sup>+</sup> released from contracting muscles during supramaximal exercise repeated three times, which is known to be associated with a large efflux of K<sup>+</sup> and lactate into plasma.*

**Material/Methods:** *Nineteen healthy students of physical education volunteered for the study. All participants performed 30s Wingate Anaerobic Test three times with 7 min rest break. During the exercise test indices of anaerobic capacity were assessed. Parameters characterizing acid base status and potassium ion concentration in plasma, whole blood and erythrocytes were measured before the exercise test, during each break between bouts and 15 min after the end of the exercise.*

**Results:** *The result showed that plasma and intraerythrocyte ion potassium concentrations were significantly increased after each bout of exercise, but exercise induced plasma hiperkalemia was levelled during the first 5 min of recovery. Simultaneously, ion potassium concentration increased in erythrocytes, but not in urine. Post exercise potassium excretion to urine was lower during 24 hours of recovery than before the exercise.*

**Conclusions:** *Obtained results suggest that erythrocytes take part in rapid changes of blood potassium level after extreme exercise. We also considered that the rapid decrease in exercise – elevated K<sup>+</sup> concentration in plasma due to their transport to erythrocytes as a prevention of the loss of potassium ion by the renal system.*

Word count: 1959

Tables: 1

Figures: 7

References: 14

Received: May 2009

Accepted: June 2009

Published: September 2009

### Address for correspondence:

Prof. dr hab. Anna Szczęśna-Kaczmarek

Academy of Physical Education and Sport, Department of Physiology, 80-336 Gdańsk, Poland, ul. K. Górskiego 1

Phone: +48 58 554-72-97, e-mail: [annaszka@awf.gda.pl](mailto:annaszka@awf.gda.pl)

## **Background**

Intense muscle contractions causes considerable changes in the intracellular concentration of electrolytes. In working muscle cells, each action potential is associated with an influx of Na<sup>+</sup> from extracellular compartment into inwards of cells and an efflux of K<sup>+</sup> from cytoplasm to extra cellular fluid [1].

The release of K<sup>+</sup> from muscle leads to a decrease in membrane potential and it is considered as an important factor in the development of muscle fatigue. It is obvious that the restoration of Na<sup>+</sup>-K<sup>+</sup> gradients between extra- and intracellular compartments is necessary before the subsequent action of a muscle. Restoration in muscle cells of Na<sup>+</sup> and K<sup>+</sup> gradients can be done by Na<sup>+</sup>-K<sup>+</sup> pump's action. The Na<sup>+</sup>-K<sup>+</sup> pump (Na<sup>+</sup>-K<sup>+</sup> - ATP-ase) action is the major mechanism allowing accumulation of K<sup>+</sup> in cells and Na<sup>+</sup> in the extracellular fluid. But the Na<sup>+</sup>-K<sup>+</sup> pump activity requires ATP for active transportation of ions. The active removal of Na<sup>+</sup> from the cell and the active uptake of K<sup>+</sup> into cells depends on the activity of Na<sup>+</sup>-K<sup>+</sup> - ATP-ase [2, 3].

During the muscle contraction, which uses up a high rate of energy, the locally available ATP supply may be limited to the extent that ion transportation and other cellular processes demanding energy cannot continue their activity at proper rates. Due to a decreased activity of Na<sup>+</sup>-K<sup>+</sup> - ATP-ase, the rapid increase in K<sup>+</sup> concentration in plasma occurred as a response to muscular contraction. It is known that K<sup>+</sup> efflux achieves values 7–16 mmol/ g wet tissue per action potential [4]. The extracellular fluid K<sup>+</sup> concentration is normally in the range of 3.35–4.45 mmol/l. Increased concentration of plasma K<sup>+</sup> is a threat to the resting membrane potential. Under these circumstances the depolarization of the resting potential can take place, which, by itself should increase neuromuscular excitability.

It is interesting that an increased level of plasma K<sup>+</sup> is not a long-lasting phenomenon, but following continued exercise, concentration of plasma K<sup>+</sup> rapidly returns to the resting values.

In such a case, the question appears what mechanism regulates such a rapid distribution of potassium ions in blood and where they are accumulated or removed.

The purpose of this paper was to examine the fate of K<sup>+</sup> released from contracting muscles during supramaximal exercise repeated three times, which is known to be associated with a large efflux of K<sup>+</sup> and lactate into plasma.

Blood may be considered as a two-compartment system comprised of red blood cells (RBC) and plasma in a ratio about 45%: 55 % at rest. Some authors proved active transportation of K<sup>+</sup> through cells membrane of erythrocytes [5, 6].

In our work we investigated the fate of K<sup>+</sup> released from contracting muscles during supramaximal exercise, taking in consideration:

- the contribution of erythrocytes to the control of the plasma concentration of K<sup>+</sup> under exercise and postexercise conditions,
- the renal secretion of K<sup>+</sup> to urine.

## **Material and Methods**

Nineteen healthy young physically active men volunteered for the study. The mean ( $\pm$ SE) age, height and weight for the group were respectively: 22.6 $\pm$ 1.77 yr, 180.9 $\pm$ 5.8 cm and 89.2 $\pm$ 11.7 kg (Table 1). The Ethical Committee of Scientific Research at the University of Medical School approved of the experimental protocol.

All experiments were conducted in the morning. Exercise was performed on a mechanical, Monark cycloergometer with automatic counting of pedal revolution. Subjects performed a warm-up lasting 5-min. before the specific test, to promote specific physiologic and motor adaptation.

After 5-min. rest, subjects performed the specific 30-s test (Wingate anaerobic test) repeated three times with 7-min. rest between each exercise bout. During the test subjects pedalled as fast as possible against a resistance of 0.075 kp per kg body mass. The maximal speed was maintained through the 30-s period.

During the test important indices characterizing anaerobic capacity were measured. Blood samples were collected in heparinized capillary tubes at rest and immediately after each exercise bout and 15-min. after the final exercise bout. Blood samples were stored on ice for analysis of: Na<sup>+</sup>, K<sup>+</sup>, Cl<sup>-</sup> concentration in plasma and whole blood, acid-base balance parameters and hematocrite. Ions concentrations in plasma and parameters of blood acid-base balance were measured immediately after blood collection. Samples of blood were frozen and thawed three times to lyse erythrocyte cell membranes before determination of Na<sup>+</sup>, K<sup>+</sup>, Cl<sup>-</sup> concentration in whole blood. Ions concentration in plasma and lysed whole blood extracts were measured using ion selective electrodes (Ciba-Corning 644 Na<sup>+</sup> /K<sup>+</sup> Cl<sup>-</sup> Analyzer). Parameters of blood acid-base status were determined by using pH/oxygen Ciba-Corning Analyser 238.

Whole blood and plasma ion concentrations were used to calculate the ion concentration of erythrocytes [7]. Urine was collected during 24 h before the exercise and 1 h, and 24 h after the completion of the last exercise bout. Urine volume was measured and urinary potassium ion concentration was determined by using ion-selective electrode in Ciba- Corning 644 Na<sup>+</sup> /K<sup>+</sup> Cl<sup>-</sup> Analyzer.

All data are presented as means with SEM. The statistical significance of difference was ascertained using Student's *t* test for paired observations.

## Results

Tab. 1. Characteristics of the examined subjects. Changes in absolute values for peak power (PP), total work (TW) and fatigue index (FI) during three times repeated 30s Wingate Anaerobic Test

Subjects	Age [years]	Body mass [kg]	PP I [W]	PP II [W]	PP III [W]	TW I [KJ]	TW II [KJ]	TW III [KJ]	FI* I [%]	FI II [%]	FI III [%]
B.J	20	98	1144	997	870	26.93	24.85	21.90	24	20	26
C.D.	25	75	796	740	879	18.47	18.95	18.35	26	22	31
C.P.	25	75	824	799	773	19.23	19.20	17.29	26	26	37
J.T.	23	98	1188	1090	1004	26.20	22.97	19.02	29	31	44
K.Ł.	20.5	110	1225	1100	1000	26.75	24.37	20.70	30	37	19
K.A.	24	98	1094	1100	1147	21.82	21.00	19.33	38	44	23
K.M.	27	90	839	850	781	21.04	19.19	19.18	18	19	30
L.R.	23	105	1133	1105	1066	26.83	22.20	21.57	22	23	33
W.J.	23	98	1143	1100	1093	26.98	24.28	25.20	26	30	39
Sz.S.	21	123	1242	1230	1424	27.43	26.12	24.49	34	33	16
D.A.	21	93	1195	1300	809	24.08	23.90	25.27	36	39	21
Z.M.	23	97	828	825	891	19.71	20.16	19.16	24	16	25
A.M.	22	77	741	739	780	17.92	17.77	16.38	21	21	14
B.A.	21	81	890	880	837	21.11	18.90	19.10	24	25	18
J.B.	23	73.5	720	898	579	17.52	18.09	16.81	19	14	20
T.M.	22	87	702	895	813	16.82	16.86	15.46	22	18	22
K.P.	21	82	884	890	813	21.07	20.61	19.26	21	20	29
S.M.	23	77	894	889	772	21.09	20.49	18.52	23	25	25
C.M.	22	77	899	800	772	16.83	13.71	12.67	29	29	26
<b>̄X</b>	<b>22.61</b>	<b>90.24</b>	<b>967.95</b>	<b>959.32</b>	<b>900.32</b>	<b>21.99</b>	<b>20.72</b>	<b>22.61</b>	<b>25.89</b>	<b>25.89</b>	<b>25.80</b>
<b>± SD</b>	<b>1.77</b>	<b>13.76</b>	<b>188.49</b>	<b>165.01</b>	<b>187.15</b>	<b>3.84</b>	<b>3.17</b>	<b>3.25</b>	<b>5.42</b>	<b>7.93</b>	<b>7.80</b>

**Performance.** During the repeated 30s Wingate anaerobic test values of Peak Power decreased from 967.95 W in the first bout to 959.2 W and 900.32 W in the second and the third bouts of exercise, respectively (Table 1). The same effect could be seen in the amount of total work. Values of total work were reduced from 21.99 KJ in the first bout to 20.72 KJ in the second repetition of the Wingate Test (Table 1).

**Potassium ions concentrations:** K<sup>+</sup> concentration in whole blood increased significantly from 72.33±7.54 mmol/l at rest to 80.27±6.12 mmol/l after three repetitions of the Wingate Test. Changes were statistically significant (p=0.007) (Fig. 1). After 15 min of recovery, the whole blood K<sup>+</sup> concentration decreased almost to the rest values.

Plasma K<sup>+</sup> concentration increased from 4.75 mmol/l at rest to 6.26 mmol/l after the first bout, but during 7 min recovery returned to normal. At the end of the second exercise bout and the third one plasma K<sup>+</sup> concentration remained significantly elevated comparing with resting values (Fig. 2).




Fig. 1. Changes in blood potassium ion concentration induced by exercise test




Fig. 2. Changes in plasma potassium ion during exercise test and recovery




Fig. 3. Erythrocyte potassium ion concentration correlated with blood potassium ion concentration

Potassium concentration in erythrocytes increased during exercise and is correlated with potassium ion concentration in blood (p=0.00002. r=0.8139) (Fig. 3). During 15 min of recovery, arising the increment of the loss of erythrocytes potassium ion was correlated with the increment of

values of blood BE and  $\text{HCO}_3^-$  concentration (Fig 4, 5). The recovery lasting fifteen minutes was sufficient to lower potassium ion concentration to the pre-exercise level.

**Blood acid-base balance.** Repeated supramaximal exercise caused changes in values of blood acid-base status. A significant decrease in blood pH was observed from  $7.40 \pm 0.01$  at rest to  $7.21 \pm 0.04$ ,  $7.15 \pm 0.05$ ,  $7.13 \pm 0.04$  at the end of the first, the second and the third exercise bouts, respectively. Also, significant changes in blood buffers were shown in the figures (Fig. 6, 7).




Fig. 4. The relationship between an increment of erythrocyte  $\text{K}^+$  concentration during recovery and blood  $\text{HCO}_3^-$  concentration.  $r = -0.4548$ ;  $p = 0.0505$




Fig. 5. The relationship between an increment of erythrocyte  $\text{K}^+$  concentration during recovery and blood BE concentration.  $r = 0.4711$ ;  $p = 0.0417$




Fig. 6. Exercise induced changes in blood  $\text{HCO}_3^-$  concentration




Fig. 7. Exercise induced changes in blood BE values

**Potassium ion concentration in urine:** Potassium ion concentration measured in urine collected during 1h and 24 h recovery after the exercise test was lower than in urine at rest. The concentration of urine potassium was  $67.16 \pm 13.0$  mmol/l and  $31.23 \pm 12.53$  mmol/l at the rest and 24 h after exercise, respectively.

## Discussion/ Conclusions

The most important finding of this study was that a rapid increase in  $\text{K}^+$  concentration in plasma during repeating supramaximal exercise did not lead to elevating of urine excretion of potassium ion. It was possible due to the participation of plasma and erythrocytes (RBC) (i.e. two

compartments of blood) in lowering large changes in plasma of potassium ion concentration released from contracting muscles.

Taking into consideration the obtained results, we can notice a protection against loss of potassium from the human body.

The quantity of  $K^+$  released during contractile action depends on the intensity of the exercise and the size of the muscle groups involved in contractions [8]. Our results show that the concentration of  $K^+$  is moderated by its uptake or release by RBCs. These observations were consistent with results of other authors [9, 10]. Transportation of  $K^+$  by human RBC is possible in both directions across the plasma membrane due to three mechanisms. The primary mechanism is active transportation by the  $Na^+-K^+$  pump [10, 11] and the second one – active transportation by the  $Na^+-K^+-2Cl^-$  cotransporter, while the third one – transportation across the  $K^+$ -ATP-sensitive channel [5, 6, 10, 11]. All mechanisms are used to move ions into RBCs.

Taking in consideration the fact that conditions after the exercise were favorable to activation of  $Na^+-K^+$ -ATP-ase due to increased concentration of catecholamines, insulin, or lactic acid [3, 12, 13], this enables us to suggest the mentioned above first mechanism as a regulating  $K^+$  transportation into RBCs.

In this study we showed not only the important role of RBC, but also renal system in  $K^+$  conservation. We observed that concentration of potassium ions in urine after supramaximal exercise was lower than at rest.

Our results are consistent with results by Medbo and Sejersted [14]. It is considered that such a rapid decrease in plasma  $K^+$  concentration is possible not only due to hiding it in erythrocytes but also to returning to muscles [5]. Described in this study turnover of  $K^+$  released from contractile muscle is of practical importance.

### **Acknowledgements**

*The Author is grateful to dr Marcin Łuszczczyk for help in statistical analysis and figures preparation.*

### **References**

1. Overgaard K, Nielsen OB, Flatman JA, Clausen T. Relations between excitability and contractility in rat soleus muscle: role of the  $Na^+-K^+$  pump and  $Na^+/K^+$  gradients. *J Physiol* 1999;518:215-225.
2. Clusen T, Gissel H. Role of  $Na^+.K^+$  pumps in restoring contractility following loss of cell membrane integrity in rat skeletal muscle. *Acta Physiol Scand* 2005;185:263-271.
3. Nielsen OB, Clausen T. The  $Na^+/K^+$  pump protects muscle excitability and contractility during exercise. *Exerc Sport Sci Rev* 2000;28:159-164.
4. Sjogard G. Exercise-induced muscle fatigue. The significance of potassium. *Acta Physiol Scand Suppl* 1990;593:1-60.
5. Lindingen ML, Horn PL, Grudzień SP. Exercise-induced stimulation of  $K^+$  in human erythrocytes. *J Appl Physiol* 1999;87:2157–2167.
6. McKelvie PS, Lindinger MI, Heigenhauser GJF, Jones NL. Contribution of erythrocytes to the control of the electrolyte changes of exercise. *Can J Physiol Pharmacol* 1991;69:984–993.
7. Buono MJ, Yeager JE. Intraerythrocyte and plasma lactate concentrations during exercise in humans. *Eur J Appl Physiol Occup Physiol* 1986;55:326-329.
8. Vollestad NK, Hallen J, Sejersted OM. Effect of exercise intensity on potassium balance in muscle and blood of man. *J Physiol-London* 1994;475:359-368.
9. McKelvie PS, Lindinger MI, Jones NL, Heigenhauser GJF. Erythrocyte ion regulations across inactive muscle during leg exercise. *Can J Physiol Pharmacol* 1992;70:1625-1633.
10. Hoffman JF. Active transport of  $Na^+$  and  $K^+$  by red blood cells. In: Fanestill DD, Schulz SG, editors. *Physiology of membrane disorders*. New York: Plenum; 1986, 221-234.
11. Lindinger MI, Grudzien SP. Exercise-induced changes in plasma composition increase erythrocyte  $Na^+,K^+$ -ATPase, but not  $Na^+-K^+-2Cl^-$  cotransporter, activity to stimulate net and unidirectional  $K^+$  transport in human. *J Physiol* 2003;533:987-997.

12. Shanmuga Sundaram KR, Padmavathi C, Sujarha Acharyja N, Vidhyalakshmi KR, Vijayan VK. Exercise-induced  $\text{Na}^+/\text{K}^+$ -ATPase activities in human red cell membrane. *Exp Physiol* 1992;77:933-936.
13. Hansen AK, Clausen T, Nielsen OB. Effect of lactic acid and catecholamines on contractility in fast-twitch muscle exposed to hyperkalemia. *Am Physiol Cell Physiol* 2006;289:C104-C112.
14. Medbo J, Sejersted OM. Plasma potassium changes with high intensity exercise. *J Physiol* 1990;421:105-122.