

2011

Muscle Strength and Power of Elite Female and Male Swimmers

Krzysztof Busko

1. Department of Biomechanics, Institute of Sport; 2. Department of Anthropology, Josef Pilsudski University of Physical Education, Poland, krzbusko@poczta.onet.pl

Jan Gajewski

1. Department of Biomechanics, Institute of Sport; 2. Department of Statistics, Josef Pilsudski University of Physical Education, Poland

Follow this and additional works at: <https://dcdansk.bepress.com/journal>



Part of the [Health and Physical Education Commons](#), [Sports Medicine Commons](#), [Sports Sciences Commons](#), and the [Sports Studies Commons](#)

Recommended Citation

Busko K, Gajewski J. Muscle Strength and Power of Elite Female and Male Swimmers. *Balt J Health Phys Act.* 2011;3(1):13-18. doi: 10.2478/v10131-011-0001-9

This Article is brought to you for free and open access by Baltic Journal of Health and Physical Activity. It has been accepted for inclusion in Baltic Journal of Health and Physical Activity by an authorized editor of Baltic Journal of Health and Physical Activity.

Muscle Strength and Power of Elite Female and Male Swimmers

DOI: 10.2478/v10131-011-0001-9

Authors' Contribution:

A – Study Design
B – Data Collection
C – Statistical Analysis
D – Data Interpretation
E – Manuscript Preparation
F – Literature Search
G – Funds Collection

Krzysztof Buśko^{1,2 (A,B,C,D,E,F,G)}, **Jan Gajewski**^{1,3 (B,D,E,F)}

¹ Department of Biomechanics, Institute of Sport, Warsaw, Poland

² Department of Anthropology, Josef Pilsudski University of Physical Education, Warsaw, Poland

³ Department of Statistics, Josef Pilsudski University of Physical Education, Warsaw, Poland

Key words: maximal muscle torque, percent muscle topography, height of jump, maximal power, swimmer, gender.

Abstract

Background: *The aim of the study was to examine gender differences in the maximal power and height of rise of the body mass centre, measured in ACMJ (akimbo counter movement jump), BCMJ (bounce counter movement jump) and CMJ (counter movement jump) and maximal muscle torque in elite swimmers who have a similar training load during the season.*

Material/Methods: *Eight male and eight female elite swimmers competing at the international level took part in the study. The maximal power and height of rise of the body mass center were measured at ACMJ, BCMJ and CMJ jumps performed on a dynamometric platform. Maximal muscle torques of elbow, shoulder, knee, hip and trunk flexors and extensors were measured under static conditions on a special stand.*

Results: *Only differences in relative muscle torque of flexors of the left and the right hip, flexors and extensors of the right knee and flexors and extensors of the trunk, the sum of relative muscle torque of the trunk and relative maximal power output measured during BCMJ were not significant between the female and male competitors, while statistically significant differences were observed between the groups in all the remaining variables of physical performances.*

Conclusions: *The results of the present study demonstrate that the male swimmers had much better performance than the female swimmers of comparable physical conditions, but when the results were calculated per body mass, gender differences in some the values were insignificant.*

Word count: 2238

Tables: 5

Figures: -

References: 17

Received: February 2010

Accepted: March 2011

Published: March 2011

Corresponding Author:

Dr hab. Krzysztof Buśko

Department of Biomechanics, Institute of Sport, Trylogii 2/16, 01-982 Warsaw, Poland

Phone: +48 22 8340812 ext. 247 E-mail: krzysztof.busko@insp.waw.pl krzbusko@poczta.onet.pl

Introduction

Physical performance capacity of males and females has been a subject of various investigations. It is a commonly accepted finding that males have higher physical performance capacity than females [1,2]. In different performance tests for maximal strength and muscle elasticity, females have reached 54-82% of the values obtained by males [1,3,4,5,6]. As reported by Patterson et al. [7], men displayed significantly higher values for relative average power and jump height in squat jump (SJ) than women. A statistically significant gender difference was found in SJ and countermovement jump (CMJ) (25% and 19% respectively) in Bosco et al. [8] study. Female subjects attained significantly lower values than men in relative power outputs in both tests – six maximal exercises of lower extremities on a cycle ergometer (18%) and 6 bouts of 10 push-offs each on an inclined plane device (48%) [9]. Malina et al. [10] reported that grip strength, 2 kg ball throw and 20 m sprint do not differ between male and female jumpers. In swimmers, relative power outputs in the Wingate test for upper extremities and swimming force in semi-tethered swimming test were significantly higher (54.9 and 74.7% respectively) in males than in females [11]. Less is known about sex differences in static muscle torques and power of lower extremities in elite swimmers. On the other hand, comparisons between the sexes should be made only if the two groups have comparable duration and intensity of the training, coaching and competition.

The aim of the study was to examine gender differences in the maximal power and height of rise of the body mass centre, measured in ACMJ (akimbo counter movement jump), BCMJ (bounce counter movement jump) and CMJ (counter movement jump) and maximal muscle torque in elite swimmers who have a similar training load during the season.

Material and Methods

The study was approved by the Ethics Committee of Institute of Sport in Warsaw, Poland. All participants were informed about the study aim and methodology as well as about the possibility of immediate resignation at any time of the experiment. Subjects agreed to the above conditions in writing. Eight male and eight female elite swimmers competing at the international level took part in the study. Their mean (\pm SD) age, body height, body mass and training experience were 21.9 ± 3.2 years, 188.8 ± 6.0 cm, 82.5 ± 3.8 kg 11.3 ± 3.5 years in men and 19.4 ± 3.7 years 175.9 ± 9.0 cm, 66.4 ± 8.9 kg, 10.3 ± 4.6 years in women, respectively. Body mass and body height were significantly different between males and females.

The maximal muscle torques of ten groups of muscles: flexors and extensors of the elbow, shoulder, hip, knee and trunk were measured in static conditions [12]. During the measurement of the muscle torque of elbow flexors and extensors the subject was sitting, with his/her arm bent at the right angle and placed on an armrest, and with the trunk stabilized. The muscle torque of shoulder flexors and extensors was measured in a sitting position. The joint angles during measurements were set at 70° and 50° during extension and flexion actions, respectively. The trunk was stabilized and the chest pressed against the testing station. The measurements of muscle torque of knee flexors and extensors were carried out on subjects in a sitting position. The hip and knee joints were bent at 90° . The subjects were stabilized at the level of anterior iliac spines and thighs, with the upper limbs crossed on his chest. The subjects were lying face down during the measurement of the muscle torque of hip extensors, and face up during the measurement of the muscle torque of hip flexors. The hip joint angle remained at 90° during both measurements. The maximal extension of the elbow, knee and hip joints was accepted as 0° . For the shoulder joint, the positioning of the arm along the trunk's side was taken as 0° . The axis of rotation during the muscle torque measurement corresponded to the axis of rotation of the torque meter.

The right arm and the left arm muscles were measured separately, always in the flexion–extension sequence. Each subject was supposed to achieve the maximal strength during measurement. The results were related to the subject's body mass and as percentage topography to the sum of muscle torque of the ten muscle groups.

The power output of lower extremities and the height of rise of the body mass center during vertical jumps were measured on a dynamometric platform with a Kistler amplifier for counter-

movement jumps (CMJ), akimbo counter-movement jumps (ACMJ) and bounce counter-movement jumps (BCMJ). The amplifier was connected to a PC via an a/d converter. The MVJ v. 3.4. software package was used for measurement. In the applied physical model the subject's body mass bouncing on the platform was reduced to a particle affected by the vertical components of external forces: the body's gravity force and the vertical component of the platform's reactive force. The maximal power and maximal height of rise of the body mass center (h) were calculated from the registered reactive force of the platform [13]. Each subject performed nine vertical jumps with maximal force on the dynamometric platform: three counter-movement jumps (CMJ), three akimbo counter-movement jumps (ACMJ) and three bounce counter-movement jumps (BCMJ). There were 5 s breaks between the CMJs and ACMJs, and 1 min breaks between the BCMJ. The jump with the highest elevation of the body's center of mass was chosen for statistical analysis.

All measurements were performed in the morning. The subjects were informed about the aims and methodology of the tests prior to the measurements.

The gender differences were calculated by formula:

$$\text{Gender differences [\%]} = ((\text{Male} * 100)/\text{Female}) - 100$$

ANOVA/MANOVA procedures with post-hoc Scheffé test were employed for comparison of means. The level of statistical significance was set at $p < 0.05$. All calculations were made with the aid of Statistica (v. 5.5, StatSoft) software package.

Results

Mean (\pm SD) values of absolute and relative maximal muscle torques of elbow, shoulder, knee, hip and trunk flexors and extensors are presented in Table 1-4. Female swimmers exhibited significantly lower torque values than male swimmers in all variables. When the values of muscle torque were calculated per unit of the body weight the gender, the difference was insignificant in flexors of the left and right hip, flexors and extensors of the right knee and flexors and extensors of the trunk. Female swimmers had a similar muscle percent topography as men with the exception of the left (2.09 ± 0.22 , 2.57 ± 0.28 N·m·kg⁻¹ respectively) and the right (2.19 ± 0.26 , 2.63 ± 0.39 N·m·kg⁻¹ respectively) extensors of the shoulder, left extensors of the elbow (1.40 ± 0.28 , 1.69 ± 0.23 N·m·kg⁻¹ respectively) and sum of relative muscle torque of the left upper extremities (Table 4).

Mean (\pm SD) values of the maximal power (Pmax) and height of rise of the body mass center measured at ACMJ, BCMJ and CMJ jumps performed on a dynamometric platform are presented in Table 5. Female swimmers generated significantly lower absolute and relative power than men with the exception of the relative maximal power measured during BCMJ. The gender difference was observed in height of all jumps, too.

Tab. 1. Mean values (\pm SD) of the maximal muscle torques [N·m] of the flexors (F) and extensors (E) of the elbow, shoulder, hip, knee and trunk

| Variables | | Female | Male | Diff. [%] | Female | Male | Diff. [%] |
|-----------|---|------------------|-------------------------------|--------------|------------------|-------------------------------|--------------|
| | | Left | Left | | Right | Right | |
| Elbow | F | 45.9 \pm 5.5 | 76.6 \pm 7.7 ^a | 66.9 | 46.0 \pm 5.0 | 75.9 \pm 11.2 ^a | 65.0 |
| | E | 30.1 \pm 4.9 | 54.6 \pm 5.2 ^a | 81.4 | 30.9 \pm 2.6 | 54.5 \pm 5.9 ^a | 76.4 |
| Shoulder | F | 38.0 \pm 7.5 | 62.8 \pm 4.1 ^a | 65.3 | 43.4 \pm 10.5 | 65.9 \pm 6.9 ^a | 51.8 |
| | E | 45.5 \pm 7.7 | 83.3 \pm 5.1 ^a | 83.1 | 48.1 \pm 10.8 | 85.5 \pm 11.6 ^a | 77.8 |
| Hip | F | 78.1 \pm 12.1 | 103.1 \pm 6.0 ^a | 32.0 | 73.3 \pm 11.3 | 103.6 \pm 9.8 ^a | 41.3 |
| | E | 335.6 \pm 73.3 | 494.4 \pm 60.8 ^a | 47.3 | 332.6 \pm 71.0 | 479.5 \pm 51.4 ^a | 44.2 |
| Knee | F | 97.0 \pm 20.5 | 139.3 \pm 12.2 ^a | 43.6 | 99.4 \pm 21.0 | 140.8 \pm 32.2 ^a | 41.7 |
| | E | 179.6 \pm 46.9 | 299.1 \pm 82.8 ^a | 66.5 | 188.1 \pm 54.6 | 291.8 \pm 72.7 ^a | 55.1 |
| Trunk | F | 137.0 \pm 29.8 | 199.8 \pm 32.8 ^a | 45.8 | | | |
| | E | 347.1 \pm 90.9 | 459.3 \pm 64.3 ^a | 32.3 | | | |

Diff. – percent gender differences, [%]; ^a - significantly different from the female, $p < 0.05$.

Tab. 2. Mean values (\pm SD) of the relative maximal muscle torques [$\text{N}\cdot\text{m}\cdot\text{kg}^{-1}$] of the flexors (F) and extensors (E) of the elbow, shoulder, hip, knee and trunk

| Variables | | Female | | Male | Diff. | Female | | Male | Diff. |
|-----------|---|-----------------|------------------------------|------|-----------------|------------------------------|-------|------|-------|
| | | Left | Right | | | Left | Right | | |
| Elbow | F | 0.69 \pm 0.05 | 0.93 \pm 0.11 ^a | 34.8 | 0.70 \pm 0.03 | 0.92 \pm 0.17 ^a | 31.4 | | |
| | E | 0.46 \pm 0.08 | 0.66 \pm 0.08 ^a | 43.5 | 0.47 \pm 0.08 | 0.66 \pm 0.09 ^a | 40.4 | | |
| Shoulder | F | 0.58 \pm 0.10 | 0.76 \pm 0.07 ^a | 31.0 | 0.66 \pm 0.15 | 0.80 \pm 0.10 ^a | 21.2 | | |
| | E | 0.69 \pm 0.07 | 1.01 \pm 0.08 ^a | 46.4 | 0.73 \pm 0.13 | 1.04 \pm 0.18 ^a | 42.5 | | |
| Hip | F | 1.19 \pm 0.18 | 1.25 \pm 0.12 | 5.0 | 1.11 \pm 0.16 | 1.26 \pm 0.14 | 13.5 | | |
| | E | 5.03 \pm 0.70 | 6.0 \pm 0.80 ^a | 19.3 | 5.0 \pm 0.73 | 5.82 \pm 0.65 ^a | 16.4 | | |
| Knee | F | 1.46 \pm 0.24 | 1.69 \pm 0.16 ^a | 15.8 | 1.50 \pm 0.28 | 1.70 \pm 0.36 | 13.3 | | |
| | E | 2.67 \pm 0.46 | 3.63 \pm 1.0 ^a | 36.0 | 2.81 \pm 0.65 | 3.53 \pm 0.86 | 25.6 | | |
| Trunk | F | 2.06 \pm 0.33 | 2.42 \pm 0.34 | 17.5 | | | | | |
| | E | 5.21 \pm 1.16 | 5.57 \pm 0.79 | 6.9 | | | | | |

Diff. – percent gender differences, [%]; ^a - significantly different from the female, $p < 0.05$.

Tab. 3. Mean values (\pm SD) of the sums of the maximal muscle torque of the right (R) and left (L) upper extremity (SUE), lower extremity (SLE), trunk (ST) and all ten muscle groups (TOTAL)

| Variables | Muscle torque [$\text{N}\cdot\text{m}$] | | Diff. |
|-----------|---|---------------------------------|-------|
| | Female | Male | |
| SUER | 168.4 \pm 24.1 | 281.8 \pm 26.2 ^a | 67.3 |
| SUEL | 159.5 \pm 20.7 | 277.3 \pm 13.6 ^a | 73.8 |
| SLER | 693.4 \pm 134.8 | 1015.6 \pm 138.4 ^a | 46.5 |
| SLEL | 690.4 \pm 136.4 | 1035.9 \pm 149.6 ^a | 50.0 |
| ST | 484.1 \pm 113.8 | 659.0 \pm 90.2 ^a | 36.1 |
| TOTAL | 2195.8 \pm 408.0 | 3269.5 \pm 329.5 ^a | 48.9 |

Diff. – percent gender differences, [%]; ^a - significantly different from the female, $p < 0.05$

Tab. 4. Mean values (\pm SD) of the sums of the relative maximal muscle torque and topography of the maximal muscle torque of the right (R) and left (L) upper extremity (SUE), lower extremity (SLE), trunk (ST) and all ten muscle groups (TOTAL)

| Variables | Muscle torque [$\text{N}\cdot\text{m}\cdot\text{kg}^{-1}$] | | | Muscle topography [%] | |
|-----------|--|-------------------------------|-------|-----------------------|------------------------------|
| | Female | Male | Diff. | Female | Male |
| SUER | 2.55 \pm 0.31 | 3.43 \pm 0.44 ^a | 34.5 | 7.77 \pm 0.89 | 8.68 \pm 1.07 |
| SUEL | 2.42 \pm 0.22 | 3.37 \pm 0.27 ^a | 39.3 | 7.38 \pm 0.91 | 8.54 \pm 0.84 ^a |
| SLER | 10.42 \pm 1.28 | 12.31 \pm 1.60 ^a | 18.1 | 31.54 \pm 0.77 | 30.99 \pm 1.56 |
| SLEL | 10.35 \pm 1.12 | 12.57 \pm 1.88 ^a | 21.5 | 31.40 \pm 1.22 | 31.60 \pm 2.26 |
| ST | 7.27 \pm 1.37 | 7.99 \pm 1.04 | 9.9 | 21.92 \pm 2.27 | 20.18 \pm 2.30 |
| TOTAL | 33.0 \pm 3.80 | 39.67 \pm 4.1 ^a | 20.2 | | |

Diff. – percent gender differences, [%]; ^a - significantly different from the female, $p < 0.05$

Tab. 5. Mean values (\pm SD) of the height of rise of the body's center of mass (h), maximal power output (P_{max}), relative maximal power output (P_{max}/mass) during akimbo counter-movement jumps (ACMJ), counter-movement jumps (CMJ) and bounce counter-movement jumps (BCMJ) on a dynamometric platform

| Variables | | P _{max} [W] | P _{max} · mass ⁻¹ [W · kg ⁻¹] | h [m] |
|-----------|--------|----------------------------|---|------------------------|
| ACMJ | Female | 1146.5±209.29 | 16.80±2.31 | 0.27±0.04 |
| | Male | 1948.0± 420.9 ^a | 23.16±5.22 ^a | 0.38±0.05 ^a |
| | Diff. | 69.9 | 37.9 | 40.7 |
| CMJ | Female | 1408.1±285.9 | 20.70±3.43 | 0.32±0.04 |
| | Male | 2318.3±464.0 ^a | 27.68±5.47 ^a | 0.42±0.05 ^a |
| | Diff. | 64.6 | 33.7 | 31.3 |
| BCMJ | Female | 1945.4±718.8 | 29.31±12.35 | 0.36±0.03 |
| | Male | 3226.0±824.1 ^a | 38.36±9.53 | 0.51±0.07 ^a |
| | Diff. | 65.8 | 30.9 | 41.7 |

Diff. – percent gender differences, [%]; ^a - significantly different from the female, $p < 0.05$.

Discussion

The results of the present study demonstrate clearly that the male subjects had much better performance than the female subjects of comparable physical conditions. As has been pointed out by Komi and Karlsson [5] the strength performance of female subjects is 20-30% lower than that of males. A similar gender difference was found by Trzaskoma and Trzaskoma [14] for relative muscle torque of the flexors and extensors of the knee and hip joints in elite athletes. On the other hand, the major contributing factor taking into account sex differences in force production is the greater muscle mass in males. When Komi and Karlsson [5] calculated the total leg force per unit of body weight, they observed very similar values for males (34.0) and females (33.8). In the present data of muscle torque, differences between the sexes were big (32.3-83.1%). The differences were smaller (6.9-46.4%) when the muscle torque was then related to the body mass, but there was a significant difference of the data, too, with the exception of the sum of relative muscle torque of the trunk, relative muscle torque of flexors of the left and right hip, flexors and extensors of the right knee and flexors and extensors of the trunk. Fidelus and Skorupski [15] suggested that in senior athletes there are specific sport-related topographies of muscle torques. On the other hand, in Buśko's [16] study absolute and relative muscle torque and percent muscle topography were significantly different between female and male elite basketball players of comparable physical conditions. In the present investigation, female swimmers had a similar muscle percent topography as men with the only exception of the left and right extensors of the shoulder, and left extensors of the elbow.

In different performance tests for maximal strength and muscle elasticity, females have reached 54-82% of values obtained by males [1,4,5,9,11]. Sienkiewicz-Dianzenza et al. [9] reported female-to-male ratio for 6 bouts of 10 push-offs each on an inclined plane device equal 48%. Komi and Bosco [1], Hudson and Owen [4] and Nelson and Martin [17] reported female-to-male ratio for height of CMJ equal 58%, 82% and 68%, respectively. In Hudson and Owen [4] paper, the difference data of height of rise of the body mass centre, velocity and use of stored elastic energy were not statistically significant between females and males of comparable physical conditions. In our study female-to-male ratio in CMJ was 31.3%. We observed significant gender differences for the value of absolute and relative power and height of jumps. Only relative power generated in BCMJ was not statistically significant between female and male swimmers of comparable physical conditions. The female-to-male ratio for all variables were not statistically significant between three different jumps.

Conclusions

The results of the present study demonstrate that male swimmers had much better performance than female swimmers of comparable physical conditions, but when the results were calculated per body mass, the gender difference in some of the values were insignificant.

References

1. Komi PV, Bosco C. Utilization of stored elastic energy in leg extensor muscles by men and woman. *Med Sci Sports* 1978;10(4):261-265.
2. Komi PV. Fundamental performance characteristics in females and males. *Med Sci Sports* 1981;14:102-108.
3. Häkkinen K. Force production characteristics of leg extensor, trunk flexor and extensor muscles in male and female basketball players. *J Sport Med Phys Fit* 1991;31(3):325-331.
4. Hudson JL, Owen MG. Performance of females with respect to males: the use of stored elastic energy. In: Winter D, Norman R, Wells R, Hayes K, Patla A, eds. *Biomechanics IX-A*. Champaign, Illinois: Human Kinetics Publishers; 1985, 50-54.
5. Komi PV, Karlsson J. Skeletal muscle fibre types, enzyme activities and physical performance in young males and females. *Acta Physiol Scand* 1978;103:210-218.
6. Miller AEJ, MacDougall JD, Tarnopolsky MA, Sale DG. Gender differences in strength and muscle fiber characteristics. *Eur J Appl Physiol* 1993;66:254-262.
7. Patterson C, Raschner C, Platzer HP. Power variables and bilateral force differences during unloaded and loaded squat jumps in high performance alpine ski racers. *J Strength Cond Res* 2009;23(3):779-787.
8. Bosco C, Tsarpela O, Foti C, et al. Mechanical behaviour of leg extensor muscles in male and female sprinters. *Biol Sport* 2002;19:189-202.
9. Sienkiewicz-Dianzenza E, Tomaszewski T, Iwańska D, Stupnicki R. Anaerobic endurance of young untrained male and female subjects. *Biomedical Human Kinetics* 2009;1:16-19. DOI: 10.2478/v10101-009-0005-y
10. Malina RM, Sławińska T, Ignasiak Z, et al. Sex differences in growth and performance of track and field athletes 11-15 years. *J Hum Kinetics* 2010;24:79-85.
11. Ogonowska A, Hübner-Woźniak E, Kosmol A, Gromisz W. Anaerobic capacity of upper extremity muscles of male and female swimmers. *Biomedical Human Kinetics* 2009;1:79-82. DOI: 10.2478/v10101-009-0020-z
12. Buśko K, Nowak A. Changes of maximal muscle torque and maximal power output of lower extremities in male judoists during training. *Hum Movement* 2008;9(2):111-115.
13. Buśko K. An attempt at the evaluation of the lower extremities power during a vertical jump on a dynamometric platform. *Biol Sport* 1988;5(3):219-225.
14. Trzaskoma Z, Trzaskoma Ł. Structure of maximal muscle strength of lower extremities in highly experienced athletes. *Physical Education and Sport* 2006;50:73-78.
15. Fidelus K, Skorupski L. Wielkość momentów sił mięśniowych w poszczególnych stawach u zawodników różnych dyscyplin sportowych [in Polish] [The levels of muscle torques in joints of athletes from various sports]. In: *Symposium teorii techniki sportowej Warszawa 26-28.11.1968 [Symposium on the Theory of the Sports Technique]*. Warszawa; 1970, 128-140.
16. Buśko K. Topografia momentów sił kobiet i mężczyzn [in Polish] [Topography of muscle torques in men and women]. *Biol Sport* 1997;14(Suppl. 7):32-36.
17. Nelson RC, Martin PE. Effects of gender and load on vertical jump performance. In: Winter D, Norman R, Wells R, Hayes K, Patla A, eds. *Biomechanics IX-B*. Champaign, Illinois: Human Kinetics Publishers; 1985, 429-433.