

2015

The importance of functional diagnostics in preventing and rehabilitating gymnast injuries with the assistance of the tensiomyography (TMG) method: A case study

Almir Atikovic

Faculty of Physical Education and Sport, University of Tuzla, Tuzla, Bosnia and Herzegovina,
almir.atikovic@untz.ba

Mitija Samardzija Pavletic

Applied Kinesiology, University of Primorska, Koper, Slovenia

Muhamed Tabakovic

Faculty of Sport and Physical Education, University of Sarajevo, Sarajevo, Bosnia and Herzegovina

Follow this and additional works at: <https://dcdansk.bepress.com/journal>



Part of the [Health and Physical Education Commons](#), [Sports Medicine Commons](#), [Sports Sciences Commons](#), and the [Sports Studies Commons](#)

Recommended Citation

Atikovic A, Pavletic MS, Tabakovic M. The importance of functional diagnostics in preventing and rehabilitating gymnast injuries with the assistance of the tensiomyography (TMG) method: A case study. *Balt J Health Phys Act.* 2015; 7(4): 29-36. doi: 10.29359/BJHPA.07.4.03

This Article is brought to you for free and open access by Baltic Journal of Health and Physical Activity. It has been accepted for inclusion in Baltic Journal of Health and Physical Activity by an authorized editor of Baltic Journal of Health and Physical Activity.

The importance of functional diagnostics in preventing and rehabilitating gymnast injuries with the assistance of the tensiomyography (TMG) method: a case study

Authors' Contribution:

- A Study Design
- B Data Collection
- C Statistical Analysis
- D Data Interpretation
- E Manuscript Preparation
- F Literature Search
- G Funds Collection

Almir Atiković^{1 ABCDEFG}, Mitija Samardžija Pavletić^{2 ABDEG},
Muhamed Tabaković^{3 ABDEG}

¹ Faculty of Physical Education and Sport, University of Tuzla, Tuzla, Bosnia and Herzegovina

² Applied Kinesiology, University of Primorska, Koper, Slovenia

³ Faculty of Sport and Physical Education, University of Sarajevo, Sarajevo, Bosnia and Herzegovina

abstract

Background The tensiomyography assessment offers information, in the time domain, regarding the following parameters: maximal radial deformation or displacement of the muscle belly, contraction time, reaction time, sustain time and relaxation time. The aim of this study is to provide information about muscle stiffness, the mechanic and contractile properties using the TMG muscles after 4 months rehabilitation process gymnast.

Material/Methods Four muscles were chosen on both lateral sides involved in artistic gymnastics performance: biceps femoris, erector spinae, gluteus maximus, rectus femoris. The testing sample in this study was taken from the Croatian Republic's senior representative who won third place 2012 and eighth place 2015 in the floor routine at the ECh.

Results The testing and measuring took place after the subject injured the lumbar region of the spinal cord and after a four-month prevention exercise program. After the first two stages of measuring, the differences can be found in: BF: -7%; ES: +17%; GM: -8%; RF: +11%. Generally speaking, a dependent t-test did not reveal significant differences in between the first and second measurement point ($t = 1.941$, $df = 39$, $P < 0.059$).

Conclusions This approach can be used to investigate top athletes who are in the process of training for muscle recovery as a result of skeletal muscle injury.

Key words skeletal muscle, contraction time, lateral symmetry, tensiomyography, men's artistic gymnastics, muscle injury

article details

Article statistics **Word count:** 2,566; **Tables:** 1; **Figures:** 1; **References:** 30

Received: October 2015; **Accepted:** November 2015; **Published:** December 2015

Full-text PDF: <http://www.balticsportscience.com>

Copyright © Gdansk University of Physical Education and Sport, Poland

Indexation: AGRO, Celdes, CNKI Scholar (China National Knowledge Infrastructure), CNPIEC, De Gruyter - IBR (International Bibliography of Reviews of Scholarly Literature in the Humanities and Social Sciences), De Gruyter - IBZ (International Bibliography of Periodical Literature in the Humanities and Social Sciences), DOAJ, EBSCO - Central & Eastern European Academic Source, EBSCO - SPORTDiscus, EBSCO Discovery Service, Google Scholar, Index Copernicus, J-Gate, Naviga (Softweco, Primo Central (ExLibris), ProQuest - Family Health, ProQuest - Health & Medical Complete, ProQuest - Illustrata: Health Sciences, ProQuest - Nursing & Allied Health Source, Summon (Serials Solutions/ProQuest, TDOne (TDNet), Ulrich's Periodicals Directory/ulrichsweb, WorldCat (OCLC)

Funding: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of interest: Authors have declared that no competing interest exists.

Corresponding author: Assistant Professor Almir Atiković, Ph.D.; University of Tuzla, Faculty of Physical Education and Sport; 2. Oktobra 1, 75000 Tuzla, Bosnia and Herzegovina; Phone/fax: +387 (0)35 278-536; E-mail: almir.atikovic@untz.ba

Open Access License: This is an open access article distributed under the terms of the Creative Commons Attribution-Non-commercial 4.0 International (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non-commercial and is otherwise in compliance with the license.

INTRODUCTION

The tensiomyography method, or in short TMG, was invented in 1983 by a group of experts from various fields at the Faculty for Electrical Engineering in Ljubljana (Slovenia), in cooperation with many institutions which were the method's early users and implementers [1]. The method has been improved since then through many prototypes and developed to the stage when commercial application of the method was possible. Although it was initially intended for medical use, the TMG method was also introduced in sports in 1996. The method was introduced by Professor Vojko Valenčič [2] and evaluated with histochemical results [3, 4], muscle force and torque [5] and EMG [6]. Skeletal muscles' velocity of the contraction is beside muscle force another very important parameter and speaks about the quality of the muscle tissue. As such it is very difficult to detect because of the subjective standardization of the measurement protocol and muscle specifics that could affect the results. To avoid those impediments, a new TMG methodology has been developed and evaluated during the last few years [7].

TMG is a non-invasive method for measuring the contractile properties of skeletal muscles. It requires no effort on the part of the subject being assessed and provides rapid, accurate information, without disrupting the daily working lives of sportsmen and women. It is used to assess muscle contraction characteristics and the contractile capacity of the superficial muscles [8, 9, 10, 11]. TMG measures geometrical changes (radial enlargement) taking place in the muscle belly during isometric contraction in response to an electrical stimulus. The assessment is made using a pressure sensor connected to a high-precision digital displacement transducer placed perpendicular to the muscle belly. The muscle is stimulated with single twitch stimulus (pulse of 1-ms duration) using two self-adhesive electrodes (2-5 cm apart) placed symmetrically to the sensor, in such a way so as not to affect the tendons [12].

The contractile properties of muscles are measured with a digital radial displacement sensor. The sensor converts physical displacement to electric impulses. These are then transported through a cable to a device that adds up the impulses and sends them to a computer with a 1 kHz frequency. The sensor response is linear with negligible noise. The dynamic response of the sensor is also very high, allowing us to distinguish differences in reaction between the fast and the slow muscle fibres. The mechanical sensor components are of the highest quality. The pressure and the resistance of the displacement rod, when placed on the muscle, does not significantly restrain the movement of the muscle.

A special sensor is placed on the muscle we wish to measure - the sensor is designed to register the muscle contraction. The muscle contraction is induced artificially with an electro stimulator, and consequently its middle part (muscle belly) enlarges. With a displacement sensor the radial enlargement of the muscle belly is measured. During measurement, the displacement sensor is pressed to the skin above the measured muscle belly radial to the muscle surface. Measuring results are presented as time/displacement curves. The newly developed report is able to provide an immediate reflection of the results obtained during the test. This report designates: optimal 90-100%, average 80-89% (monitor), and poor (concern) <80% muscle responses and muscle symmetries. In this study we will present just a few possibilities of

the methods' application in men's artistic gymnastics: analyses of lateral and functional symmetry, muscle adaptation to a specific sport or exercise, muscle potentiation and fatigue and muscle recovery from injury. The main aim of this study is to provide information about muscle stiffness, the mechanic and contractile properties of the skeletal muscles using the TMG with high level gymnast from men's artistic gymnastics as well as to demonstrate the usefulness of this method to evaluate the muscles after rehabilitation process.

MATERIAL AND METHODS

SUBJECT

The testing sample in this survey was taken from the Croatian Republic's senior representative in gymnastics, Andrej Korosteljev, born on December 15th 1996 in Osijek (Croatia). In 2012 in Montpellier in France, he qualified for the finals at the European Juniors Championships and won the third place in the floor routine and the eighth place for seniors three years later in 2015. The subject's morphological characteristics are: body height 172 cm, body weight 63 kg and the body mass index 21.3 kg/m². In the past six months he has trained on average for 4 hours 5 times a week (twice a day). The mechanism of the lumbar region of spinal cord injury occurred after a bad landing caused by a mat slip during his training. The injury diagnosis was given by a specialist physician after receiving the results of the magnetic resonance imaging (MRI). The injury was followed by a prevention program which lasted from the 6th of June 2014 till the 24th of October 2014. All the measuring and testing was conducted in the Center for Sports Medicine, Ljubljana, Slovenia [ID: 15121995].

The participant of this study was fully informed of the potential risks associated with the study and signed written consent forms previously approved by the Research Ethics Committee of the University of Tuzla in line with the criteria of the Helsinki Declaration for research involving human beings.

ELECTRICAL STIMULATION

Two circular (5 or 3 cm in diameter depending on the muscle belly size), self-adhesive electrodes (Axelgaard, Pulse) were placed bipolarly on the skin above the muscle belly. The diameter of the electrodes was chosen on the basis of the muscle size to avoid activation of the synergistic muscles or co-activation. Electrodes were placed 5 cm medially and laterally from the measuring point. A single rectangular monophasic pulse of 1 ms duration from an electro-stimulator (TMG-S1, Furlan & Co. ltd.) was delivered to the electrodes, transcutaneously eliciting a twitch muscle contraction.

SIGNAL RECORDINGS

The TMG signals were stored on a portable PC. The digital TMG signal was directly sampled with the Matlab Compiler Toolbox using a sampling frequency of 1 kHz. Two maximal responses were stored and averaged for further analyses. The maximal stimulation amplitude was identified as the minimal stimulation amplitude needed for the response with the highest displacement amplitude (Dm) [13].

Four time parameters as a measure for contractile dynamics were calculated from the response: Delay time (Td) as the time between the electrical impulse and 10% of the contraction; Contraction time (Tc) as the time between 10% and 90% of the contraction; Sustain time (Ts) as the time between 50% of the contraction and 50% of relaxation; Relaxation time (Tr) as the time between 90% and 50% of relaxation. These parameters are used to assess the stiffness of the muscle and its balance in relation to other muscle structures, muscle chains (flexion-extension) and extremities (right and left) [2].

MEASURED MUSCLES

Four muscles were chosen on both lateral sides involved in artistic gymnastics performance: Biceps Femoris (BF), Erector Spinae (ES), Gluteus Maximus (GM), Rectus Femoris (RM).

STATISTICAL ANALYSIS

SPSS 21.0 was used for statistical analysis, which included descriptive statistics (means and standard deviations), correlational analyses, t-tests. Paired t-test was performed to determine whether there were any significant differences after the recovery program. The significant level was defined as ($P < 0.05$).

RESULTS

General Biceps Femoris (BF) muscle LS is very high, 95%. General Erector Spinae (ES) muscle LS is slightly lower than recommended, 76%. LS of the contraction time is significantly lower than recommended. The muscle on the right side is slower. We recommend activation exercises for the right side. General Gluteus Maximus (GT) muscle LS is very high, 92%. The muscle is slightly faster than the reference value. General Rectus Femoris (RF) muscle LS is slightly lower than recommended, 79%. LS of maximal displacement is significantly lower than recommended. It is lower on the left side. The muscle is significantly faster than the reference value. We recommend strength exercises for both sides. The relatively good bilateral symmetry in delay times (Td) between muscles indicates that neural transmission to the muscles is not compromised by the disc protrusion. Bilateral symmetry in the rectus femoris is low; the left rectus femoris is slower/weaker. Bilateral symmetry in the erector spinae is also low; here there could be 2 scenarios. First: The right erector spinae is slow/weak, which - combined with the weak/slow left RF - could be the origin of the low back pain problems. In this case the pelvis would be rotated to the right in the transverse plane during dynamic movements. Second: An alternative scenario is that the left erector spinae is tightened (low Dm and low Tc) because of pain/herniation to the left. In our opinion we believe the first scenario is the most likely. We can conclude that there is a no statistically significant difference between the mean in the first and the second measurement point ($t=1.941$, $df=39$, $P < 0.059$).



Table 1. Statistical parameters TMG for Tc, Ts, Tr, Dm, Sim - Lateral Symmetry

Parameters	DM	Muscle/Side			Muscle/Side			Muscle/Side			Muscle/Side		
		BF		R-L	ES		R-L	GM		R-L	RF		R-L
		Right	Left	%	Right	Left	%	Right	%	%	Right	Left	%
Tc [ms]	1	29.78	30.61	2.1	18.58	15.03	19.10	40.91	37.46	8.43	19.79	21.62	8.51
	2	24.35	28.30	13.95	15.84	15.49	2.20	34.15	39.97	14.56	18.24	19.86	8.15
Ts [ms]	1	218.00	185.65	14.83	169.99	52.97	68.83	79.02	65.45	17.17	188.66	59.33	68.55
	2	233.10	224.62	3.63	89.00	49.46	44.42	91.26	169.41	46.13	30.04	41.28	27.22
Tr [ms]	1	69.38	52.00	25.0	147.40	33.26	77.43	32.44	25.14	22.50	166.15	32.03	80.72
	2	87.28	55.73	36.14	71.33	32.44	54.52	50.31	127.05	60.40	10.84	18.56	41.59
Dm [mm]	1	8.03	8.74	8.12	4.98	3.90	21.68	9.83	9.86	0.30	7.64	4.21	44.89
	2	9.63	8.34	13.39	7.45	6.99	6.17	14.74	14.58	1.08	6.98	6.09	12.75
Td [ms]	1	24.35	25.53	4.62	21.95	18.09	17.58	52.47	47.13	10.17	22.14	21.37	3.47
	2	24.33	23.31	4.19	19.41	19.51	0.51	47.05	36.47	22.48	21.42	21.75	1.51
Sim [%]	1	95			76			92			79		
	2	88			93			84			90		



Legend: (DM) Date of first measurement 06.06.2014; second measurement 24.10.2014.

1st measurement

ES Erector Spinae



ES Erector Spinae



2nd measurement

Legend: ms - time in milliseconds; mm - displacement in mm

Fig. 1. Erector Spinae (ES) at the 1st and the 2nd measurement - the right and the left side comparison

DISCUSSION

The main aim of this study is to provide information about muscle stiffness, the mechanic and contractile properties of the skeletal muscles using the TMG with high level gymnast from men's artistic gymnastics as well as to demonstrate the usefulness of this method to evaluate the muscles after rehabilitation process.

From a general point of view the picture is much better. The parameters which were weak in the first measurement ES and GM testing have increased and improved their values. In addition, other muscles have also improved their contraction time which is now faster than it was after the first measuring. Measuring has also shown improvement during the period between two measurements. The improvement itself is visible in the stabilization of parameters which can affect the appearance of the injury. The risk is now satisfactory and significantly decreased. The TMG method can be used as a further contribution to optimizing the process of rehabilitation and physical recovery of athletes with muscle injuries. When athletes get injured, they, the team, the coach, and the organization, all suffer. TMG assessments could be useful in identifying and selecting young athletes. A simple methodology is an important factor for the feasibility of the method. In addition, it is objective, non-invasive, rapid and selective, and can give repetitive information.

In the available publications, many authors describe skeletal muscles. Zagorc et al. [13] made a comparison of contractile parameters among twelve skeletal muscles of inter-dance couples. From the sample of 8 top dancers in Slovenia (average age: 19.1 ± 3.6 years). Tc is for BF R: 34.2 ± 7.7 , L: 34.3 ± 9.1 , ES R: 18.3 ± 2.2 , L: 17.5 ± 2.2 , GM R: 23.5 ± 2.5 , L: 23.2 ± 1.6 , RM R: 32.9 ± 5.8 , L: 35.5 ± 6.5 . García-García's study [14], the sample was composed of ten professional road cyclists (age 27.5 ± 5.5 years old; height 178.2 ± 7.8 cm; body mass 65.6 ± 5.4 kg) who had planned the "Vuelta a España" as the main competition of the season. Tc for RF: 45.9 ± 16.2 and BF: 28.2 ± 5.2 . Samardžija Pavlečić et al. [15] focused on the contractility comparison between 10 sample muscles of ($n = 26$) Slovenian gymnasts. The tests showed similar results in three muscles BF: 25.0 ± 9.0 ; ES: 14.8 ± 1.7 ; RF: 21.1 ± 3.2 . According to internet source and results of Spanish football club Rayo Vallecano, [16] based on a case study of 1 top male football player who plays center position (age: 18 years, height 172 cm, weight 68 kg) for the junior team Rayo Vallecano in Spain, a comparison of contractile parameters was made among four skeletal muscles. Tc for BF R: 22.19, L: 22.11, Sim (94%), RF R: 20.38, L: 22.85, Sim (80%). In a study on athletes in sprint discipline [17] (age: men 25 years, height 173 cm, weight 79 kg). Tc for BF R: 19.65, L: 27.37, Sim (76%), ES R: 11.81, L: 12.51, Sim (93%), GM R: 27.73, L: 27.35, Sim (97%), RM R: 22.75, L: 24.41, Sim (89%). In Loturco et al. study [18] on forty-one high-level track and field athletes from power ($n = 22$; age 27.2 ± 3.6 years; height 180.2 ± 5.4 cm; weight 79.4 ± 8.6 kg) and endurance events (endurance runners and triathletes; $n = 19$; age 27.1 ± 6.9 years; height 169.6 ± 9.8 cm; weight 62.2 ± 13.1 kg) had the mechanical properties of their rectus femoris (RF) and biceps femoris (BF) assessed by TMG. Tc BF = 14.3 ± 2.3 vs. 19.4 ± 3.3 ms, Tc RF = 18.3 ± 2.8 vs. 22.9 ± 4.0 ms.

Floor exercises in men's artistic gymnastics is one of the most technically complex and the most difficult events, requiring a gymnast to have adequate physical fitness skeletal muscle of the lower extremities and low back. Most

injuries in artistic gymnastics are related to floor landing [19, 20, 21, 22, 23, 24, 25]. This phase not only affects the gymnast's final rank during competition [26], but also entails a high risk of injury, mainly due to the high impact magnitudes of 14 to 18 body weight applied to one leg [19], and to the mat's instability [27, 28]. Landing imposes forces on the body that must be absorbed primarily by the musculoskeletal components of the lower extremities. If the loads become too great for the body to accommodate, a potential injury situation arises [29, 30]. Designing programs with exercises for prevention and good landing in gymnastics, separated from the whole routine, would help to decrease impact forces during landing and reduce the possibility of injury.

CONCLUSION

TMG results can help the coach to establish the future training load and this is very important for prevent muscle injuries. Coaches should monitor such muscle adaptation more often at different training or competition periods during the year. All of these facts suggest that organised and strict work should be undertaken to maintain health, reduce the risk of injuries and ensure proper body development in artistic gymnastics.

Limitation of this study is that the sample size was small and other limitation is this relatively new method of testing skeletal muscle and for now there is no large number of studies for comparison with other similar studies. We believe that this TMG method, after training program of rehabilitation can be a useful device for muscle mechanic diagnostics.

REFERENCES

- [1] Dias P, Fort J, Marinho D, Santos A, Marques M. Tensiomyography in physical rehabilitation of high level athletes. *The Open Sports Sciences Journal*. 2010;3:47-48.
- [2] Valenčić V. Direct measurement of the skeletal muscle tonus. In: Popovic D, editor. *Advances in External Control of Human Extremities*. Beograd: Nauka; 1990.
- [3] Dahmane R, Knez N, Valenčić V, Erzen I. Tensiomyography, a non-invasive method reflecting the percentage of slow muscle fiber in human skeletal muscles. In: *Book of Abstract: Life Sciences September 28th to October 1st 2000, Gozd Martuljek Slovenia; 2000, 29.*
- [4] Dahmane R, Valenčić V, Knez N, Erzen I. Evaluation of the ability to make non-invasive estimation of muscle contractile properties on the basis of the muscle belly response. *Med Biol Eng Comput*. 2000;38:51-55.
- [5] Šimunič B. Modelling of longitudinal and transversal skeletal muscle belly deformation. Doctor Thesis. Ljubljana: Faculty of Electrical Engineering; 2003.
- [6] Kerševan K. Comparison of biomechanical and myoelectrical biceps brachii response in men. Master Thesis. Ljubljana Faculty of Electrical Engineering; 2002.
- [7] Šimunič B, Rozman S, Pišot R. Detecting the velocity of the muscle contraction. In: *Proceedings Book of 3rd International Scientific Conference "New Technologies in Sports - NTS"*. Sarajevo; 2005.
- [8] Valenčić V, Knez N. Measuring of the skeletal muscles dynamic properties. *Artific Org*. 1997;21:240-242.
- [9] Valenčić V, Djordjević S, Knez N, et al. Contractile properties of skeletal muscles detection by tensiomyographic measurement method. In: *2000 Pre-Olympic Congress, Abstract 507*. Brisbane, Australia; 2000.
- [10] Valenčić V, Knez N, Šimunič B. Tensiomyography: detection of skeletal muscle response by means of radial muscle belly displacement. *Biomedical Engineering*. 2001;1:1-10.
- [11] Dahmane R, Djordjević S, Smerdu V. Adaptive potential of human biceps femoris muscle demonstrated by histochemical, immunohistochemical and mechanomyographical methods. *Med Biol Eng Comput*. 2006;44(11):999-1006.
- [12] Rodríguez-Matoso D, Rodríguez-Ruiz D, Sarmiento S, Vaamonde D, Da Silva-Grigoletto ME, García-Manso JM. Reproducibility of muscle response measurements using tensiomyography in a range of positions. *Revista Andaluza de Medicina del Deporte (Centro Andaluz de Medicina del Deporte Sevilla, España)*. 2010;3(3):81-86.

- [13] Zagorc M, Šimunić B, Pišot R, Oreb G. A comparison of contractile parameters among twelve skeletal muscles of inter-dance couples. *Kinesiologia Slovenica*. 2010;16(3):57-65.
- [14] García-García O. Relación entre parámetros de tensiomiografía y potenciales indicadores del rendimiento en ciclistas profesionales [The relationship between parameters of tensiomyography and potential performance indicators in professional cyclists]. *Revista Internacional de Medicina y Ciencias de la Actividad Física y el Deporte*, 2013;13(52):771-781. Spanish.
- [15] Samardžija Pavletić M, Kolar, E., Šimunić B. Tensiomyography in artistic and rhythmic gymnastics. In: *Proceedings Book of 2nd International Scientific Conference "Gymnastics Federation of Slovenia"*, 23.1.2015. Portorož, 63-65.
- [16] Rayo Vallecana Juvenil DH\AR2011072687 [Available at <http://g-se.com/uploads/biblioteca/ar2011072687.pdf>][Accessed on 1 March, 2015].
- [17] Buckley M. Gaining Muscle Performance Insight with Tensiomyography TMG North America™ ("TMG") - Freelap USA [<http://www.freelapusa.com/gaining-muscle-performance-insight-with-tensiomyography/>][Accessed on 1 March, 2015].
- [18] Loturco I, Gil S, Frota de Souza Laurino C, et al. Differences in muscle mechanical properties between elite power and endurance athletes: a comparative study. *J Strength Cond Res*. 2014 Dec 24.
- [19] Panzer VP. Lower Extremity Loads in Landings of Elite Gymnasts. (Doctoral dissertation), Oregon: University of Oregon; 1987.
- [20] Lindner KJ, Caine DJ. Injury patterns of female competitive club gymnasts. *Can J Sport Sci*. 1990;15:254-261.
- [21] Meeusen R, Borms J. Gymnastic injuries. *Sports Med*. 1992;13:337-356.
- [22] McNitt-Gray JL, Munkasy BA, Welch M, Heino J. External reaction forces experienced by the lower extremities during the take-off and landing of tumbling skills. *Technique*. 1994;14:10-16.
- [23] Kirialanis P, Malliou P, Beneka A, Gourgoulis V, Giofstidou A, Godolias G. Injuries in artistic gymnastic elite adolescent male and female athletes. *J Back Musculoskeletal Rehabil*. 2002;16(4):145-151.
- [24] Kirialanis P, Dallas G, Di Cagno A, Fiorilli G. knee injuries at landing and take-off phase in gymnastics. *Sci Gymnastics J*. 2015;7(1):17-25.
- [25] Marinšek M. Basic landing characteristics and their application in artistic gymnastics. *Sci Gymnastics J*. 2010;2(2):59-67.
- [26] Leskošek B, Čuk I, Karacsony I, Pajek J, Bučar M. Reliability and validity of judging in men's artistic gymnastics at the 2009 University Games. *Sci Gymnastics J*. 2010;2(1):25-34.
- [27] Arampatzis A, Brüggemann GP, Klapsing GM. A three dimensional lower leg-foot model to determine the influence of various gymnastic mats on foot during landings. *Med Sci Sport Exer*. 2002;34:130-138.
- [28] Arampatzis A, Klapsing GM, Brüggemann GP. The effect of falling height on muscle activity and foot motion during landings. *J Electromyogr Kines*. 2003;13:533-544.
- [29] Dufek JS, Bates BT. The evaluation and prediction of impact forces during landing. *Med Sci Sport Exer*. 1990;22(3):370-377.
- [30] Čuk I, Marinšek M. Landing quality in artistic gymnastics is related to landing symmetry. *Biol Sport*. 2013;30(1):29-33. Epub 2013 Jan 21.

Cite this article as:

Atiković A, Samardžija Pavletić M, Tabaković M. The importance of functional diagnostics in preventing and rehabilitating gymnast injuries with the assistance of the tensiomyography (TMG) method: a case study. *Balt J Health Phys Act*. 2015;7(4):29-36.