

2016

Healthy aging: Quality of life and health behaviours in students of the University of the Third Age – A preliminary study

Justyna Krzepota

*Department of Physical Culture and Health Promotion, University of Szczecin, Poland,
justyna.krzepota@univ.szczecin.pl*

Dorota Sadowska

Department of Physiology, Institute of Sport-National Research Institute, Warsaw, Poland

Beata Florkiewicz

Department of Physical Culture and Health Promotion, University of Szczecin, Poland

Follow this and additional works at: <https://dcdansk.bepress.com/journal>



Part of the [Health and Physical Education Commons](#), [Sports Medicine Commons](#), [Sports Sciences Commons](#), and the [Sports Studies Commons](#)

Recommended Citation

Juškelienė V, Kalinkevičienė A. Health disorders and exercising behaviour that characterize the transitional stage of first-year students at the Lithuanian University of Educational Sciences . Balt J Health Phys Act. 2016; 8(4): 117-126. doi: 10.29359/BJHPA.08.4.13

This Article is brought to you for free and open access by Baltic Journal of Health and Physical Activity. It has been accepted for inclusion in Baltic Journal of Health and Physical Activity by an authorized editor of Baltic Journal of Health and Physical Activity.

Healthy aging: Quality of life and health behaviours in students of the University of the Third Age - A preliminary study

Authors' Contribution:

- A Study Design
- B Data Collection
- C Statistical Analysis
- D Data Interpretation
- E Manuscript Preparation
- F Literature Search
- G Funds Collection

Justyna Krzepota^{1 ACDEF}, Dorota Sadowska^{2 CDEF}, Beata Florkiewicz^{1 BF}

¹ Department of Physical Culture and Health Promotion, University of Szczecin, Poland

² Department of Physiology, Institute of Sport-National Research Institute, Warsaw, Poland

abstract

Background The aim of the presented study was to analyse self-assessment of the quality of life and general health behaviours among the students of the University of the Third Age.

Material/Methods The research was conducted on a sample of 81 persons (M = 62.6, SD = 6.1). The tool used for assessing the quality of life was the Quality of Life Questionnaire (WHOQoL-BREF, Polish version), and the methodological basis for investigating health behaviours was the Health Behaviour Inventory.

Results The analysis conducted among all participants showed statistically significant positive correlations between the General Index of Intensity of Health Behaviours (GIIHB) and the quality of life in the Social Relationships domain ($R_s = 0.23$; $p \leq 0.05$). Statistically significant positive correlations were also found between Positive Thinking in the Health Behaviour Inventory and two WHOQoL-BREF domains: the Psychological domain ($R_s = 0.24$; $p \leq 0.05$) and the Social Relationships domain ($R_s = 0.26$; $p \leq 0.05$).

Conclusions The results of our research may prove very helpful in designing health promotion programs for an ageing society. Further longitudinal studies with a larger sample size are needed in order to identify more clearly the relationships that occur between health behaviours and the quality of life, especially in the process of ageing.

Key words Quality of life, health behaviors, University of the Third Age, elderly, aging

article details

Article statistics **Word count:** 3,022; **Tables:** 3; **Figures:** 0; **References:** 36

Received: Februar 2016; **Accepted:** October 2016; **Published:** December 2016

Full-text PDF: <http://www.balticsportscience.com>

Copyright © Gdansk University of Physical Education and Sport, Poland

Indexation: AGRO, Celdes, CNKI Scholar (China National Knowledge Infrastructure), CNPIEC, De Gruyter - IBR (International Bibliography of Reviews of Scholarly Literature in the Humanities and Social Sciences), De Gruyter - IBZ (International Bibliography of Periodical Literature in the Humanities and Social Sciences), DOAJ, EBSCO - Central & Eastern European Academic Source, EBSCO - SPORTDiscus, EBSCO Discovery Service, Google Scholar, Index Copernicus, J-Gate, Naviga (Softweco, Primo Central (ExLibris), ProQuest - Family Health, ProQuest - Health & Medical Complete, ProQuest - Illustrata: Health Sciences, ProQuest - Nursing & Allied Health Source, Summon (Serials Solutions/ProQuest, TDOne (TDNet), Ulrich's Periodicals Directory/ulrichsweb, WorldCat (OCLC)

Funding: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of interest: Authors have declared that no competing interest exists.

Corresponding author: Dr Justyna Krzepota; Department of Physical Culture and Health Promotion, University of Szczecin, al. Piastów 40b, blok 6, 71-065 Szczecin, Poland; tel.: +48 91 444 2789; fax: 91 444 2734; e-mail: justyna.krzepota@univ.szczecin.pl

Open Access License: This is an open access article distributed under the terms of the Creative Commons Attribution-Non-commercial 4.0 International (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non-commercial and is otherwise in compliance with the license.

INTRODUCTION

Due to a steady rise in life expectancy and an increased number of elderly people, there is a need to undertake measures aimed at improving the health-related quality of life of senior citizens. Health-related quality of life and well-being for all individuals, as well as improving the health, function, and quality of life especially of older adults are a new topic area in the Healthy People 2020 Project [1]. In order to take the correct preventive action related to the quality of life of older people, it is necessary to identify the factors determining their lifestyle and their influence on well-being [2]. Among the factors influencing the quality of life, researchers often emphasize the importance of physical activity [3], nutritional habits [4, 5], BMI [6, 7] time devoted to sleep [8], and mental aspects [9]. Another related research topic concerned the importance of adult life behaviours and their influence on the quality of life in older age [10]. Also the predisposing factors for depression in older people were researched [11]. International research conducted among elderly people in 22 countries showed that the most important quality of life characteristics were having energy, feeling happy, having well-functioning senses, and being free from pain [12]. According to scholars, despite evidence concerning the influence of health behaviours on health, there is still too little data on the relationship between health behaviours and the quality of life, especially in older people [13]. For example, it has been shown that smokers exhibit a lower level of the health-related quality of life [14]; they also complain about difficulties in performing daily activities and loss of well-being significantly more often than non-smokers [15]. Research indicates that being overweight [6], lacking in sleep [8], depression [16] and social isolation [17] are all factors lowering the quality of life in older age.

Among the factors influencing the quality of life in older age, health behaviours and practices promoting health and well-being seem to be of special importance [13]. Health education programs conducted among people over 60 years of age contribute to a better understanding and an improvement in health behaviours of the participants [18]. Moreover, research indicates that access to late-life learning, in the form of attending the University of the Third Age, also has a positive influence on the quality of life of older people [19]. Apart from fulfilling the need for intellectual development, the University of the Third Age gives an opportunity for social integration and making new acquaintances, at the same time providing a pleasurable way of spending free time in older age [20]. Authors emphasize that participation in classes for a period longer than one semester is related to a higher level of satisfaction with life and greater psychological adjustment [21], and it reduces depression levels [22]. Nevertheless, introducing changes in health behaviours of older people is not an easy task and it entails a number of challenges [23]. Therefore, a clear understanding of which health behaviours influence the general quality of life and its specific domains is crucial for designing health promotion programs for older people.

AIM

The aim of the presented study was self-assessment concerning the Overall Quality of Life and General Health, as well as the quality of life in four domains: Physical Health, Psychological, Social Relationships, and Environmental among students of the University of the Third Age, and their general health behaviours

such as: Proper Nutritional Habits, Preventive Behaviours, Positive Thinking and Health Practices.

MATERIAL AND METHOD

PARTICIPANTS

The research was conducted on a sample of 81 persons ($M = 62.6$, $SD = 6.1$), including 67 women and 14 men. All the participants were students of the University of the Third Age from the West Pomeranian region in Poland. Respondents filled out the questionnaire voluntarily. The respondents were divided into two groups based on Spirduso's classification [24], where the dividing mark is the age of 65. The group under 65 years of age involved 55 participants ($M = 59.2$, $SD = 3.5$), while the group over 65 years of age consisted of 26 participants ($M = 69.6$, $SD = 4.0$). The local Bioethical Committee approved of the research project.

PROCEDURES

The tool used for assessing the quality of life was a Polish version of the abridged World Health Organization Quality of Life Questionnaire (WHOQoL-BREF) developed by Wołowicka and Jaracz [25], and the methodological basis for investigation health behaviours was Juczyński's Health Behaviour Inventory [26].

MEASUREMENT OF THE QUALITY OF LIFE: THE POLISH VERSION OF THE WHOQOL-BREF QUESTIONNAIRE.

The WHOQoL-BREF questionnaire consists of 26 questions. The first two questions were analysed separately. They pertained to self-perception of the respondents' Overall Quality of Life and General Health. The remaining 24 questions assessed four aspects of the quality of life (Physical Health - 7 questions, Psychological - 6 questions, Social Relationships - 3 questions, and Environmental - 8 questions). The respondents were asked to mark their answers using a five-level rating scale (from 1 to 5 points). The quality of life in respective domains was expressed as the mean value, calculated according to the key and guidelines provided by the authors [25].

MEASUREMENT OF HEALTH BEHAVIOUR: HEALTH BEHAVIOUR INVENTORY

The General Index of Intensity of Health Behaviours (GIIHB) was calculated by adding the results for all the 24 statements included in the Health Behaviour Inventory. The obtained GIIHB ranged from 24-120 points. The results were then converted to standard units and interpreted in terms of the sten score system. According to the guidelines presented by the author of the questionnaire, the respondents were divided into three groups: with a high GIIHB: 7-10 stens, an average GIIHB: 5-6 stens, and a low GIIHB: 1-4 stens. Four categories of health behaviours were analysed separately: Proper Nutritional Habits, Preventive Behaviours, Health Practices and Positive Thinking.

The presented study examines four categories: Proper Nutritional Habits, which according to Juczyński [26] takes into consideration mainly the type of foods consumed (e.g. wholegrain bread, vegetables or fruit); Preventive Behaviours, which include adhering to medical recommendations and obtaining information concerning health and sickness; Health Practices, which

pertain to daily habits connected with sleep, recreation and physical activity; Positive Thinking, which means avoiding strong emotions, stress, tension and depressing situations.

They were calculated as the mean value of the results in the analysed category (ranging from 1.0-5.0), following the adopted diagnostic key. The higher the result, the healthier the habits.

STATISTICAL ANALYSES

The results were analysed statistically with the aid of STATISTICA 10.0 program. The distributions of variables were examined using the Saphiro-Wilk test. The relevance of differences was assessed with the aid of U Mann Whitney test. Relationships between the variables were analysed using Spearman's rank correlation test and a correlation coefficient was calculated for each pair of variables. The level of statistical significance was accepted at $p \leq 0.05$.

RESULTS

The analysis of results of the WHOQoL-BREF Quality of Life Questionnaire for all the participating students of the University of the Third Age ($n = 81$) indicates that the respondents assessed their Overall Quality of Life at the average level of $M = 3.64$ ($SD = 0.58$), while the mean value for General Health assessment was $M = 3.28$ ($SD = 0.81$). The University of the Third Age students evaluated their quality of life in the Physical Health domain to be $M = 12.66$ ($SD = 1.64$) and the average level for the Psychological domain was $M = 13.67$ ($SD = 1.95$). As to the Social Relationships domain and the Environmental domain, the results were $M = 13.81$ ($SD = 2.43$) and $M = 13.63$ ($SD = 1.76$), respectively.

The analysis of health behaviours using the Health Behaviour Inventory showed that the GIIHB for all the participating University of the Third Age students oscillated around $M = 86.31$ ($SD = 11.30$). A low GIIHB exhibited 17.3% of the respondents, 53.1% average and 29.6% high. The mean value for Proper Nutritional Habits was $M = 3.74$ ($SD = 0.67$) and for Preventive Behaviours it amounted to $M = 3.66$ ($SD = 0.62$). As far as Positive Thinking is concerned, the participants scored an average of $M = 3.50$ ($SD = 0.57$), while for Health Practices it was $M = 3.50$ ($SD = 0.59$).

In Table 1 the WHOQoL-BREF Quality of Life Questionnaire results for the University of the Third Age students under 65 years of age were compared with the results obtained by the group of participants over 65 years old.

Table 1. Mean and standard deviation for the WHOQoL-BREF quality of life questionnaire results in groups of University of the Third Age students under 65 and over 65 years old and the significance of differences between the groups

WHOQoL-BREF	< 65 age M (SD)	≥ 65 age M (SD)	U	p
Overall Quality of Life	3.65 (0.62)	3.62 (0.50)	0.21	0.84
General Health	3.35 (0.82)	3.15 (0.78)	0.98	0.33
WHOQoL-BREF domain				
Physical Health	12.78 (1.79)	12.40 (1.26)	0.72	0.47
Psychological	14.10 (1.90)	12.77 (1.77)	2.69	0.01**
Social Relationships	13.96 (2.60)	13.49 (2.04)	0.76	0.45
Environmental	13.70 (1.84)	13.48 (1.61)	0.47	0.64

** $p \leq 0.01$

The only statistically relevant difference between the groups was found in the Psychological domain. The group of University of the Third Age students below 65 years old assessed their quality of life in the Psychological domain (M = 14.10, SD = 1.90) statistically higher ($p \leq 0.01$) than the group over 65 years of age (M = 12.77, SD = 1.77). In all the remaining domains, age was not a differentiating factor. What is more, no statistically relevant differences between the groups were noted as far as the assessment of the Overall Quality of Life and General Health is concerned.

The Health Behaviour Inventory results for University of the Third Age students under and over 65 years old did not exhibit statistically relevant differences in the assessment of low, high and average GIIHB. Low GIIHB was attested in 21.8% of University of the Third Age students below 65 years of age and in 7% of students over 65 years of age. Average GIIHB characterized 47.3% of students below 65 years old and 65.4% of students over 65 years old. High GIIHB, in turn, was achieved by 30.9% of students below 65 years of age and 26.9% of students over 65 years of age. Comparative analysis also did not reveal any statistically significant differences between the groups in the mean values for respective categories of health behaviours (Proper Nutritional Habits, Preventive Behaviours, Positive Thinking and Health Practices, see: Table 2).

Table 2. Mean and standard deviation for the Health Behaviour Inventory results in groups of University of the Third Age students under 65 and over 65 years old, and the significance of differences between the groups

Health Behaviour Inventory	< 65 age M (SD)	≥ 65 age M (SD)	U	p
General Index of Intensity of Health Behaviours	85.76 (12.31)	87.46 (8.90)	-0.14	0.89
Proper Nutritional Habits	3.72 (0.74)	3.78 (0.49)	0.01	0.99
Preventive Behaviours	3.61 (0.66)	3.72 (0.51)	-0.58	0.56
Positive Thinking	3.52 (0.58)	3.46 (0.57)	0.88	0.38
Health Practices	3.44 (0.58)	3.62 (0.60)	-1.15	0.25

The next stage of statistical analysis focused on investigating whether there is a relationship between a subjective view of the quality of life assessed using WHOQoL-BREF and the health behaviours declared in the Health Behaviour Inventory. The analysis of correlations conducted among all the participating University of the Third Age students (see: Table 3) showed statistically relevant relationships between GIIHB and the quality of life in the domain of Social Relationships. Respondents with higher GIIHB assessed their quality of life higher in the aspect of the Social Relationships domain ($R_s = 0.23$; $p \leq 0.05$). Statistically relevant positive correlations were also found between Positive Thinking in the Health Behaviour Inventory and two WHOQoL-BREF domains, the Psychological domain ($R_s = 0.24$; $p \leq 0.05$) and the Social Relationships domain ($R_s = 0.26$; $p \leq 0.05$).

Table 3. Correlations between the results of WHOQoL-BREF and health behaviour inventory conducted among students of the University of the Third Age

WHOQoL-BREF	WHOQoL-BREF domain					
	Overall Quality of Life	General Health	Physical Health	Psychological	Social Relationships	Environmental
Health Behaviour Inventory						
General Index of Intensity of Health Behaviours	0.14	0.15	0.14	0.14	0.23* $p = 0.04$	0.16
Proper Nutritional Habits	0.11	0.12	0.08	0.00	0.13	0.02
Preventive Behaviours	0.05	0.13	0.12	0.07	0.19	0.12
Positive Thinking	0.08	0.13	0.10	0.24* $p = 0.03$	0.26* $p = 0.02$	0.15
Health Practices	0.19	0.00	0.08	0.00	0.06	0.16

* $p \leq 0.05$

DISCUSSION

In the present study, the quality of life, health behaviours and the relationship between these factors were analysed. 53.1% of the respondents obtained the General Index of Intensity of Health Behaviours (GIIHB) at a moderate level, whereas 29.6% exhibited a high level and 17.3% a low level. The results of the present study are in agreement with previous studies concerning these factors conducted among University of the Third Age students (46% moderate level, 28% high level and 26% low level, respectively) [27]. These results are slightly different from the data obtained by other authors [28] who noted the highest percentage of participating University of the Third Age students to have a high GIIHB (43.2%), 39.7% moderate and 17.1% low one. Proper health behaviours are important to prevent diseases and to reduce the death rate among people. Therefore, it is very comforting to find that in our research, as well as in the above-mentioned studies, it was the smallest percentage of respondents that was characterized by low GIIHB.

In our study, the analysis of the quality of life self-assessment results showed that the participating University of the Third Age students perceived their Overall Quality of Life as average ($M = 3.64$) and General Health was assessed by them slightly lower ($M = 3.28$). These results are similar to results obtained in studies on different Universities of the Third Age in Poland [20] and abroad [29]. It is reassuring that, as present study indicates, there is no significant difference in results between people under and over 65 years of age, which implies that despite aging, the perception of Overall Quality of Life and General Health among the University of the Third Age students participating in the research does not deteriorate.

Respondents under 65 years of age assessed their quality of life in all the four domains (Physical Health, Psychological, Social Relationships and Environmental) slightly better than those over 65 years old. Nevertheless, statistically relevant differences were noted only in the Psychological domain ($p \leq .01$). The observed difference between the two groups most probably indicates a lower level of satisfaction with life, self-satisfaction and a sense of meaning in life among the participants over 65 years of age. These observations confirm the results of other research [30] in which the influence of age on assessing the quality of life in the Psychological domain was noted in a study of older people in Turkey. Analysing changes perceived in the quality of life over age, the researchers observed that the highest score in the Psychological domain was achieved by people aged 65-69, while in the 70-74 and 75-89 age brackets, as well as in people over 80 years old, the perceived quality of life gradually decreased. Other authors [31] indicate that mental aspects play an extremely significant role in the subjective perception of the quality of life, and a lower quality of life was observed in elderly people with depression.

Moreover, one of the most important factors influencing the perception of the quality of life by older people is functional fitness. In present research both the respondents below and over 65 years of age assessed their quality of life in the Physical Health domain as the lowest. It seems understandable, since the process of ageing and past diseases cause deterioration in physical fitness, functional mobility and the ability to work, as well as increased pain and discomfort.

Earlier research results [32] indicate that Social Relationships and Environmental domains are important in the subjective perception of the quality of life by elderly people. Moreover, other authors [33] suggest that according to the elderly social and environmental domains in elderly assisted living facilities are low. Our results also showed that for the majority of respondents, the Social Relationships domain of the quality of life was significantly associated with GIIHB and one of the health behaviour categories, i.e. Positive Thinking. Also Psychological domain of the quality of life was strongly correlated with Positive Thinking. These findings are in agreement with the observations made by other authors [e.g. 34], which imply that positive thinking, positive feeling and positive emotions increase the quality of physical and psychological health, and the subjective well-being.

The limitation of the present study is that the analysis was narrowed to the health behaviour variables adopted in the questionnaire, which creates some interpretation difficulties, as few authors [35, 36] have analysed the

relationship between health behaviours and the quality of life, based on the Health Behaviour Inventory and WHOQoL-BREF, respectively. Also the specific nature of the sample (University of the Third Age students) and a limited number of participants make it impossible to draw general conclusions true for the whole population of older adults and elderly people. Nevertheless, the results of the present research are a source of important information concerning the health behaviours undertaken by the students of the University of the Third Age and provide an insight into a subjective perception of the quality of life and health among senior citizens.

CONCLUSION

In conclusion, the present study concentrated on how University of the Third Age students perceive their quality of life in relation to the health behaviours they undertake. Quality of life, especially in Social Relationship and Psychological domains, is shaped by health behaviours. Positive Thinking is important in Social Relationships and Psychological domains. Also the level of health behaviours is positively correlated with the Social Relationships domain. Further longitudinal studies with a larger sample size are needed in order to identify more clearly the relationships that occur between health behaviours and the quality of life, especially in the process of ageing. The results of such research may prove very helpful in designing health promotion programs for an ageing society.

REFERENCES

- [1] Office of Disease Prevention and Health Promotion. Healthy People. Available at <http://www.healthypeople.gov/2020/topics-objectives/topic/health-related-quality-of-life-well-being>. [Accessed on 15 November, 2014].
- [2] Pullen C, Walker SN, Fiandt K. Determinants of health-promoting lifestyle behaviors in rural older women. *Fam Community Health*. 2001;24(2):49-72.
- [3] Vagetti GC, Barbosa Filho VC, Moreira NB, Oliveira VD, Mazzardo O, Campos Wd. Association between physical activity and quality of life in the elderly: A systematic review, 2000-2012. *RBP*. 2014;36(1):76-88.
- [4] Jiménez-Redondo S, De Miguel BB, Banegas JG, Mercedes LG, Gomez-Pavon J, Vives CC. Influence of nutritional status on health-related quality of life of non-institutionalized older people. *J Nutri Health Aging*. 2014;18(4):359-364.
- [5] Alfonso-Rosa RM, del Pozo-Cruz B, del Pozo-Cruz J, del Pozo-Cruz JT, Sañudo B. The relationship between nutritional status, functional capacity, and health-related quality of life in older adults with type 2 diabetes: A pilot explanatory study. *J Nutr Health Aging*. 2013;17(4):315-321.
- [6] Lopez-Garcia E, Banegas JB, Gutierrez-Fisac JL, Pérez-Regadera AG, Díez-Gañan L, Rodriguez-Artalejo F. Relation between body weight and health-related quality of life among the elderly in Spain. *Int J Obesity*. 2003;27(6):701-709.
- [7] Bottone FG, Hawkins K, Musich S, et al. The relationship between body mass index and quality of life in community-living older adults living in the United States. *J Nutr Health Aging*. 2013;17(6):495-501.
- [8] Faubel R, Lopez-Garcia E, Guallar-Castillón P, et al. Sleep duration and health-related quality of life among older adults: a population-based cohort in Spain. *Sleep*. 2009;32(8):1059-1068.
- [9] Chan SWC, Chiu HF, Chien WT, Goggins W, Thompson D, Hong B. Predictors of change in health-related quality of life among older people with depression: a longitudinal study. *Int Psychogeriatr*. 2009;21(06):1171-1179.
- [10] Holahan CK, Suzuki R. Adulthood predictors of health promoting behavior in later aging. *Int J Aging Hum Dev*. 2004;58(4):289-313.
- [11] Green BH, Copeland JRM, Dewey ME, et al. Risk factors for depression in elderly people: A prospective study. *Acta Psychiatr Scand*. 1992;86(3):213-217.
- [12] Molzahn A, Skevington SM, Kalfoss M, Makaroff KS. The importance of facets of quality of life to older adults: An international investigation. *Qual Life Res*. 2010;19(2): 293-298.
- [13] Mo PKH, Mak WWS. The influence of health promoting practices on the quality of life of community adults in Hong Kong. *Soc Indic Res*. 2010;95(3):503-517.

- [14] Holahan CK, Holahan CJ, North RJ, Hayes RB, Powers DA, Ockene JK. Smoking status, physical health-related quality of life, and mortality in middle-aged and older women. *Nicotine Tob Res.* 2013;15(3):662-669.
- [15] Laaksonen M, Rahkonen O, Martikainen P, Karvonen S, Lahelma E. Smoking and SF-36 health functioning. *Prev Med.* 2006;42(3):206-209.
- [16] Lin JH, Huang MW, Wang DW, et al. Late-life depression and quality of life in a geriatric evaluation and management unit: an exploratory study. *BMC Geriatr.* 2014;18(14):77, doi:10.1186/1471-2318-14-77.
- [17] Hawton A, Green C, Dickens AP, et al. The impact of social isolation on the health status and health-related quality of life of older people. *Qual Life Res.* 2011;20(1):57-67.
- [18] Huang LH, Chen SW, Yu YP, Chen PR, Lin YC. The effectiveness of health promotion education programs for community elderly. *J Nurs Res.* 2002;10(4):261-270.
- [19] Hebestreit L. The role of the University of the Third Age in meeting needs of adult learners in Victoria, Australia. *Aust J Adult Learn.* 2008;48(3):547-565.
- [20] Zielińska-Więczkowska H, Kędziora-Kornatowska K, Ciemnoczołowski W. Evaluation of quality of life (QoL) of students of the University of Third Age (U3A) on the basis of socio-demographic factors and health status. *Arch Gerontol Geriatr.* 2011;53(2):e198-e202.
- [21] Ordonez TN, Lima-Silva TB, Cachioni M. Subjective and psychological well-being of students of a university of the third age. *Dement Neuropsychol.* 2011;5(3):216-225.
- [22] Ordonez TN, Batistoni SS, Cachioni M. [Depressive symptoms in older adults studying at a third age open university]. *Rev Esp Geriatr Gerontol.* 2011;46(5):250-255. Spanish.
- [23] Burton LC, Paglia MJ, German PS, Shapiro S, Damiano AM. The effect among older persons of a general preventive visit on three health behaviors: smoking, excessive alcohol drinking, and sedentary lifestyle. The Medicare Preventive Services Research Team. *Prev Med.* 1995;24(5):492-497.
- [24] Spirduso WW. *Physical Dimensions of Aging.* Champaign, Ill: Human Kinetics; 1995.
- [25] Wołowicka L, Jaracz K. Polska wersja WHOQOL - WHOQOL 100 i WHOQOL BREF [The Polish versions of WHOQOL 100 and WHOQOL Bref]. In: Wołowicka L, editor. *Jakość życia w naukach medycznych [The quality of life in medical sciences].* Poznań: Wydawnictwo Akademii Medycznej; 2001, 231-281. Polish.
- [26] Juczyński Z. Inwentarz Zachowań Zdrowotnych - IZZ [Health Behaviour Inventory - HBI]. In: Juczyński Z, editor. *Narzędzia pomiaru w promocji i psychologii zdrowia [Measurement tools in the promotion and health psychology].* Warszawa: Pracownia Testów Psychologicznych; 2001, 116-122. Polish.
- [27] Kozieł D, Kaczmarczyk M, Naszydlowska E, Gałuszka R. Wpływ kształcenia w Uniwersytecie Trzeciego Wieku na zachowania zdrowotne ludzi starszych [The influence of studying at the third age university on elderly people's health behaviour]. *Studia Medyczne.* 2008;12:23-28. Polish.
- [28] Smoleń E, Gazdowicz L, Żyłka-Reut A. Zachowania zdrowotne osób starszych [Health behaviors in elderly population]. *Pielęgniarstwo XXI Wieku.* 2011;36(3):5-9. Polish.
- [29] Carvalho-Loures M, Celeno-Porto C, Alves-Barbosa M, Freire-Filha LG. Women's quality of life: University of the Third Age, Goiás, Brazil. *Rev Salud Publica.* 2010;12(3):391-401.
- [30] Arslantas D, Ünsal A, Metintas S, Koc F, Arslantas A. Life quality and daily life activities of elderly people in rural areas, Eskişehir (Turkey). *Arch Gerontol Geriatr.* 2009;48(2):127-131.
- [31] Chan SW, Chiu HF, Chien WT, Thompson DR, Lam L. Quality of life in Chinese elderly people with depression. *Int J Geriatr Psychiatry.* 2006;21(4):312-118.
- [32] García EL, Banegas JR, Pérez-Regadera AG, Cabrera RH, Rodríguez-Artalejo F. Social network and health-related quality of life in older adults: a population-based study in Spain. *Qual Life Res.* 2005;14(2):511-520.
- [33] Bodur S, Cingil DD. Using WHOQOL-BREF to evaluate quality of life among Turkish elders in different residential environments. *J Nutr Health Aging.* 2014;13(7):652-656.
- [34] Fredrickson BL. The role of positive emotions in positive psychology: The broaden-and-build theory of positive emotions. *Am Psychol.* 2001;56(3):218-226.
- [35] Dębska G, Korbel-Pawlas M, Zięba M, Ławska W, Lubierda A. Jakość życia a zachowania zdrowotne w grupie uczestników Uniwersytetu Trzeciego Wieku - badanie wstępne [Quality of life vs. health behaviours among respondents attending the University of the Third Age (U3A) - a preliminary study]. *Zdrowie Publiczne.* 2012;122(1):48-54. Polish.
- [36] Kurpas D, Mroczek B, Jasinska M, Bielska D, Nitsch-Osuch A, Kassolik K. Health behaviors and quality of life among patients with chronic respiratory disease. *Adv Exp Med Biol.* 2013;788:401-406.

Cite this article as:

Krzepota J, Sadowska D, Florkiewicz B. Healthy aging: Quality of life and health behaviours in students of the University of the Third Age - A preliminary study. *Balt J Health Phys Act.* 2016;8(4):127-135.